

Language acquisition in developmental disorders

Michael S. C. Thomas

School of Psychology, Birkbeck College, London

To appear in: In M. Kail, M. Hickmann & M. Fayol (Eds.), *Proceedings of the International Conference on First and Second Language Acquisition*, Paris 2006.

Address for correspondence:

Dr. Michael Thomas
Developmental Neurocognition Laboratory
School of Psychology
Birkbeck College, University of London
Malet Street, Bloomsbury
London WC1E 7HX, UK
Email: m.thomas@bbk.ac.uk
Tel.: +44 (0)20 7631 6386
Fax: +44 (0)20 7631 6312

Keywords: Developmental disorders, Atypical development, Language disorders, Williams syndrome, Down syndrome, Autism, Specific Language Impairment, Dyslexia, Developmental delay, Computational modelling, Connectionism, Neuroconstructivism, Compensation, Redundancy, Functional brain imaging, Fractionation of language

Running head: Language acquisition in developmental disorders

1: Introduction

What light can developmental disorders shed on language development? Can they reveal the extent to which language development is channelled by biological constraints? Can they demonstrate whether language learning relies on general cognitive mechanisms or whether it is domain-specific? In this chapter, we consider what has been learned by the comparison of language development across multiple disorders, as well as the unresolved issues that still exist in this field.

First, let us clarify what is meant by developmental disorders. Developmental disorders can be split into four groups. The first are disorders caused by well-understood genetic abnormalities, such as Down syndrome (three copies of chromosome 21) and Williams syndrome (deletion of ~25 genes from one copy of chromosome 7). In these neurogenetic disorders, cognitive impairments are typically not restricted to a single cognitive domain. The second group are disorders defined on the basis of behavioural deficits, such as dyslexia, Specific Language Impairment and autism. In these disorders, behavioural genetics indicates sometimes substantial heritability, but the causal genes are not yet known and may well not be mutations (that is, they may be spectrum disorders corresponding to an unlucky accumulation of normal genetic variations that each add a small risk for the target disorder). In these disorders, it is sometimes argued that the deficits are restricted to single cognitive domains (e.g., reading in dyslexia, language in Specific Language Impairment) but there remain doubts as to whether these disorders are indeed homogeneous rather than behavioural clusters with milder associated deficits and heterogeneous causes. The third group are disorders where there is learning disability but its cause is unknown. The final group are disorders caused by environmental factors, such as viral infections or an impoverished environment, be it cognitive or biological (such as in Foetal

Alcohol syndrome). The first and last of these four groups index the primary locus of causality – the first group nature, the last group nurture – while the middle two reflect our current lack of knowledge. A given behavioural impairment may be generated in more than one way. For example, poor reading may be the consequence of either dyslexia or limited opportunities to learn to read. Our discussion will predominantly focus on the first two groups – neurogenetic and behavioural disorders (see, e.g., Goldin-Meadow, 2005, for discussion of language development under conditions of impoverished input). Disorders of development that are caused by early acquired brain damage will be considered briefly in section 4.

Both dissociation and association methodologies have been applied to developmental disorders of language (see Bishop, 1997, Karmiloff-Smith, 1998, Temple, 1997, for discussion). Where *ability A* develops normally but *ability B* develops atypically, a possible inference is that they are subserved by independent systems that do not interact during development. Where *ability A* and *ability B* both develop atypically, one possible inference is that a common system subserves their development; another is that they are subserved by two systems that causally interact across development (Morton, 2004). Very different explanatory frameworks have been deployed in interpreting language deficits in developmental disorders. On the one hand, some researchers have extended the logic of adult cognitive neuropsychology to developmental disorders, hypothesising that patterns of behavioural deficits should be related to normal modular theories of the language system (for the appropriate age); deficits are then viewed as the failure of individual components to develop (e.g., Clahsen & Temple, 2003). On the other hand, other researchers stress the interactive, adaptive nature of the developmental process, arguing that the normal adult modular structure is the product of the developmental

process rather than a precursor to it and, since components interact across development, impairments are likely to spread; moreover, genetic effects in disorders are typically widespread in the brain rather than equivalent to focal lesions; together, these researchers infer that the language system in developmental disorders may be qualitatively atypical and therefore have no direct correspondence to the normal language system (e.g., the *neuroconstructivist* position; see Karmiloff-Smith, 1998; Mareschal et al., 2007; Thomas & Karmiloff-Smith, 2002, 2005). Currently, then, some researchers believe that at best, developmental disorders of language may help reveal the structure of the normal language system while others argue that at worst, disorders may tell us nothing about normal language development. One goal of this chapter is to indicate where in between these two extremes the truth may lie.

The following examples illustrate the types of claims that have been made about language development in development disorders. It has been argued that Specific Language Impairment may be a genetic failure of language (and in some cases, only syntax) to develop against a background of otherwise normally developing cognition (e.g., as assessed by non-verbal intelligence tests) (Pinker, 1999; van der Lely, 2004). Williams syndrome, a rare neurogenetic disorder, shows an uneven cognitive profile, with relatively strong language ability (for overall mental age), a particular weakness in visuospatial construction and a background of learning disability. Based on early reports of this disorder, Pinker (1994, 1999) argued that language might develop normally in this disorder despite deficits in general cognition. In high-functioning individuals with autism, it has been argued that the structural parts of language can be acquired appropriately but these individuals do not master its use in social situations (something rather crucial for language) (Happé, 1994). These claims revolve around disorders that exhibit dissociations, but equally we need

explanations of associations, for example where all aspects of language development are delayed in a disorder but individuals nevertheless seems to follow normal milestones, though perhaps terminating at a lower level of sophistication (e.g., as in Down syndrome). What property of a cognitive system could produce general language delay? Speculations about how language development can go wrong rely on a detailed understanding of how it works in the normal case.

2. Language as a learning problem

The effects of developmental damage to the language system may be quite different to the effects of acquired damage in adulthood, because in the former case one cannot assume that there is already a language system in place (Thomas & Karmiloff-Smith, 2002). Instead, developmental deficits must be interpreted as disruptions to an adaptive learning process. Theories of language development differ depending on how tightly constrained they view the learning process to be (very tightly in nativist theories, where environmental input serves to ‘trigger’ adult states; weakly in empiricist theories where structure in input-output mappings serves to construct the adult state from more general resources). Minimally, developmental disorders must be viewed in terms of changes to the constraints under which language development takes place, whether learning is tightly or loosely constrained. But learning theories bring into play a range of other concepts. These include the interactions between different information sources or processing mechanisms, the importance of the quality of input and output representations, changes in plasticity with age, compensation between processing components when some are initially impaired, and the possibility of redundancy (i.e., multiple developmental pathways to success).

At the most abstract level, Tager-Flusberg and Sullivan (1998) characterised normal language development as involving the integration of three streams of information, about the physical world, about people, and about the structure of language itself. Ultimately, these will form the basis of lexical semantics, pragmatics, and phonology/syntax respectively. These information streams are depicted in Figure 1. The most important point is that language development involves the integration of these information sources – to use some linguistic structure to convey some meaning to achieve some social ends. But the integration process may be complicated: some types of information may be redundantly available in more than one information stream; or information in one stream may help resolve ambiguities in the other and so aid its acquisition (the basis of the developmental notion of *bootstrapping*). In this way, Chiat (2001) has emphasised how disorders of language development must construe observed impairments in terms of the way each disorder changes the problem of learning the mapping from sound to meaning and from meaning to sound.

=====

Insert Figure 1 about here

=====

Cross-syndrome comparisons are potentially most informative about the different ways in which the developmental process can be deflected. Figure 2 demonstrates illustrative data from our lab of the sorts of patterns that can be observed when disorders are compared (Annaz, Johnson & Thomas, 2006). These data depict cross-sectional developmental trajectories for 18 children with Williams syndrome (WS), 15 children with Down syndrome (DS), 16 high-functioning children with autism (HFA), and 17 low-functioning children with autism (LFA) between the ages of 5 and 12, against a typically developing (TD) sample of 25 children. The left panel

shows performance on a standardised test of receptive vocabulary (a task where the child has to point to the picture that goes with a word), while the right panel shows performance on a non-verbal test of visuo-spatial construction (a task where the child has to complete a simple puzzle, building a target from geometric shapes). In both cases, test (mental) age is plotted against chronological age.

Two of the disorders show similar profiles across verbal and non-verbal measures, illustrating *developmental associations*. For the HFA group, development is slightly below the TD trajectory but within the normal range, while the DS group shows very delayed and only slowly improving performance on both measures. By contrast, the WS group shows development parallel to and just below the normal range for language (similar to the HFA group), but very delayed development on visuo-spatial construction (similar to the DS group). Meanwhile, the LFA group shows poor performance on language development (indeed, there is no significant improvement with chronological age in this cross-sectional sample) but then development within the normal range for visuo-spatial construction (similar to the HFA group). These latter two cases illustrate *developmental dissociations*.

=====

Insert Figure 2 about here

=====

Such cross-syndrome comparisons have been carried out to explore associations and dissociations within the domain of language itself, both in early development (Tager-Flusberg & Sullivan, 1998) and later childhood (Fowler, 1998) (see also Rice, Warren & Betz, 2005). These comparisons focused on phonology, syntax, semantics and pragmatics, and identified several contrasting profiles. For high-functioning children with autism, problems primarily occur in pragmatics, in line

with the social disengagement typical of the disorder. For low-functioning children with autism, there are additionally problems with lexical semantics and concept formation. Problems in lexical semantics and concepts also characterise the development of children with learning disability (or ‘mental retardation’, to use US terminology). In Williams syndrome, language development is mostly characterised by delay but with a relatively successful eventual outcome. However, there are also differences in pragmatics but now the pattern is of hypersociability, with an elevated interest in using language for social engagement. In Down syndrome, problems appear to primarily impact on the structural aspects of language, especially phonology and those parts of language that rely on phonological distinctions (morphology, syntax). SLI and dyslexia are also viewed as behavioural disorders that primarily impact on structural language information, with sub-types emphasising difficulties in phonology, semantics, or syntax. The contrast between these disorders is included in Figure 1.

What kinds of conclusions have been drawn from these comparisons? Fowler (1998) noted that pragmatics and semantics appear to be most closely tied to overall mental age across different disorders, while phonology and syntax can dissociate. Either pragmatics and semantics involve more general systems, or their successful development requires interactions between a greater number of cognitive components. McDonald (1997) contrasted various populations in which language acquisition is broadly successful (including WS and HFA) with those in which language acquisition is unsuccessful (including DS and SLI, but also late L1 and L2 learners). Her conclusion was that good representations of speech sounds (phonology) are crucial in predicting eventual successful acquisition. When the individual cannot encode the basic phonological contrasts over which the rules of language operate, prognosis is

poor. However, as Morton (2004) argues, many cognitive components typically contribute to the successful development of an overall system, and if any one of these is impaired (and no redundancy is present) the system may fail to develop normally. Good phonology may be a necessary but not sufficient requirement for successful language acquisition.

Both Fowler (1998) and Tager-Flusberg and Sullivan (1998) were struck by the absence of radically different pathways by which language can be developed. In most disorders, acquisition exhibits similarities to the normal trajectory, proceeding through a common sequence and via common milestones (as far as acquisition progresses in a given disorder). Their common conclusion was that these similarities must be the result of invariant internal biological constraints that shape language development in all the disorders. Thus Fowler argued that "...language acquisition [is] heavily constrained by brain structure" (1998, p.309), while Tager-Flusberg and Sullivan concluded that "there are not multiple alternative ways of acquiring language, though as each of these components [phonology, semantics, and syntax] develops over time, they may become integrated in different ways, which lead to syndrome-specific profiles" (1998, p.231). An alternative possibility is that, on computational grounds, some of the similarities to typical development are likely to emerge from the common structure of the problem domain that these children are acquiring when they seek to communicate meaning via sound (Thomas, 2005a).

In the next two sections, we consider two more detailed examples of language acquisition in developmental disorders. These stress how important it is to view atypical language development in terms of the trajectory of an adaptive learning system operating under altered constraints (computational or informational). The first example shows how research has progressed over a decade or more of investigating

language development in Williams syndrome, and introduces the idea of *redundancy* in language development. The second example of Specific Language Impairment reveals the emergence of new methods to address key issues in the atypical development of language, and introduces the idea of *compensation*.

3. The case of language development in Williams syndrome

Williams syndrome has been much studied over the last fifteen years due to the uneven cognitive profile observed in this neurogenetic disorder (Donnai & Karmiloff-Smith, 2000). Figure 2 depicts one of the most salient dissociations observed in standardised testing: a disparity between receptive vocabulary and visuospatial constructive skill. Individuals with WS also show a hypersociable or ‘over-friendly’ personality profile (Jones et al., 2000), with a relative strength in facial recognition. By contrast, they have relative weaknesses in numeracy and problem solving skills, and overall IQs typically fall between 50 and 70. Based on the early findings of Ursula Bellugi from a small number of individuals with the disorder, Pinker (1994, 1999) argued that WS might constitute a genetic dissociation in which grammar develops normally but general intelligence is impaired – in support of a wider argument that normal language development involves innate, domain-specific mechanisms. Although, as with any disorder, there is variability, individuals with WS often have a surprising facility with language compared to some of their other abilities, and compared to other disorders with comparable overall mental age such as Down syndrome (e.g., as shown in Figure 2). A dissociation of this nature encourages the idea that developmental disorders might serve to ‘fractionate’ the cognitive system into its component parts. The simple fractionation proposed by Pinker (1994) is shown in Figure 3a.

=====

Insert Figure 3a and 3b about here

=====

These initial claims inspired a burst of research on WS that has lasted fifteen years and incorporated investigation of the genetic basis of the disorder, its effects on brain development, and a detailed consideration of the cognitive abilities of these individuals using more sensitive experimental tasks. Research on brain development has tended to indicate that the genetic effects of the mutation are fairly widespread rather than focal, consistent with most neurogenetic disorders that affect cognition (Toga, Thompson & Sowell, 2006). By contrast, research on the cognitive abilities of these individuals has revealed an increasingly complex and fine-grained picture. In the domain of language, the most salient characteristic in WS is that development is delayed. Early in childhood, these children’s language ability is on a par with children with DS (Paterson et al., 1999). Only in late childhood and adolescence does WS language development stretch away, while that of DS asymptotes. In most published empirical studies, the performance of individuals with WS is compared to a typically developing control group matched for *mental age* (MA); performance is very rarely at the level of a control group matched for chronological age. MA comparisons implicitly accept that there is no dissociation between language ability and mental age in WS (although the notion of MA itself is weakened by disorders in which there is an uneven cognitive profile).

Various studies have reported dissociations within the domain of language, for instance problems in learning spatial prepositions, difficulties in the pragmatics of conversation, and problems with more complex aspects of morphology. Thomas and Karmiloff-Smith (2003) reviewed the literature at the turn of the century and

identified two types of emerging hypothesis. The *Semantics-Phonology Imbalance hypothesis* suggests that individuals with WS are relatively strong in their language development but that it occurs in a subtly different way. In WS, there might be greater emphasis on the sounds of words and less emphasis on their precise meaning. For example, in early language development, children with WS show vocabulary growth ahead of the normal markers of semantic development such as referential point and object sorting (see Thomas, 2005a, for a review). By contrast, the *Conservative hypothesis* suggests that there is nothing atypical about language development in WS – it is entirely in line with mental age (i.e., it is delayed). What anomalies there are stem from other characteristics of the disorder such as the visuospatial deficit that causes problems in learning spatial prepositions (*in, on, under*) and the hypersociable profile that leads these individuals to use language strategically in a way to capture and maintain attention in social interactions (see, e.g., Thomas et al., in press, for an example in the context of unusual vocabulary use in WS). Under the Conservative hypothesis, language in WS is made to look more impressive by comparing it to other cognitive domains in which there are particular weaknesses (e.g., visuospatial construction) and to other disorders in which there are known phonological processing problems, such as DS and SLI (e.g., Ring & Clahsen, 2005).

As research has progressed in WS, methodological problems such as restricted sample sizes and inappropriate control groups have increasingly been addressed. Brock (in press) recently reviewed the status of the two competing hypotheses. He found that the *Conservative hypothesis* has gained progressively more support over the *Imbalance hypothesis*. Delay remains the most salient feature of language development in WS and performance appears to be in line with the level of general cognition (excluding the visuospatial deficit). While there are some anomalies

compared to MA-matched control groups, most of these appear to stem from other non-verbal aspects of the disorder. One exception may be receptive vocabulary (e.g., as shown by the data in Figure 2, left panel). This skill is puzzlingly strong even compared to the rest of language and the disparity remains to be explained. Brock (in press) argues that the slow and anomalous early phase of language development in WS combined with the eventual relative success in acquisition implicates *redundancy*. That is, early language development in the disorder does not exploit the normal combination of information sources and cognitive processes; it finds a pathway to success that takes longer but is nonetheless eventually successful. This position contrasts with that of Tager-Flusberg and Sullivan (1998) who, as we saw earlier, argued against alternative pathways for successful language acquisition.

To offer a concrete example of this redundancy, Laing et al. (2002) identified deficits in shared attention in toddlers with WS. Although these toddlers scored well on dyadic interactions (sharing attention with the caregiver), they exhibited deficits in triadic interactions, where attention had to be shifted between the caregiver and an object that was being played with. The deficit was a consequence of their elevated interest in (and fixation on) the face of the caregiver. It is thought that triadic interactions are an important contributor to learning object names in situations where the caregiver labels an object that is being played with (“*Look at the ball! This is a ball!*”) (e.g., Tomasello & Farrar, 1986). The toddler with WS may therefore be deprived of this information source to some extent in their language development. However, explicit labelling is not the only route to learning object names, and while development is slower, these children do succeed in vocabulary acquisition. The inference is therefore that other redundant pathways to success are followed, which are less efficient and take longer.

Overall, research into the cognitive profile of individuals with WS has tended to produce increasingly fine-scale fractionations between different abilities even within cognitive domains. Although the initial fractionation in WS was argued to be between language and cognition as shown in Figure 3a, the current picture of is closer to that shown in Figure 3b. This fine-scaled fractionation contrasts with the coarse and widespread effect of the genetic mutation on brain development. One can make this point more starkly: in WS, the granularity of genetic differences in cortex is far coarser than the level of cognitive modules; the impact on cognitive development is a granularity of subsequent fractionations considerably finer than the level of cognitive modules (Thomas, 2006). The difference in granularity between genetic and cognitive effects arises because cognitive structure is the result of a developmental process that exaggerates or attenuates the effects of atypical constraints on learning, depending on the cognitive domain (Karmiloff-Smith, 1998). In the next section, we will see how new methods are important to specifying the nature of this developmental process.

4. The case of language development in Specific Language Impairment

SLI is a behaviourally defined disorder diagnosed by the presence of a deficit in language development in the presence of apparently normal non-verbal development and the absence of any obvious neurological impairment or environmental cause. It is a heritable disorder but the precise genes involved are unknown. SLI is sometimes conflated with the British KE family. Affected members of this family were reported to have particular problems with language and the cause was traced to a mutated FOXP2 gene on chromosome 7 (see Marcus & Fisher, 2003). As with WS and in keeping with other neurogenetic disorders, subsequent research has indicated that cognitive differences and brain differences between affected and unaffected family

members are more widespread than the domain of and substrate for language (Watkins, Dronkers & Vargha-Khadem, 2002; Watkins et al., 2002). However, behaviourally defined SLI is not caused by the FOXP2 mutation (Newbury et al., 2002).

SLI is a disorder that primarily impacts on syntax and phonology, although its particular features depend on the language being acquired. It appears to be a heterogeneous disorder, with subtypes that differentially impact morphology/syntax, semantics, and pragmatics (Bishop & Norbury, 2002). Three principal theories have been advanced for the cause of behaviourally defined SLI. First, SLI has been explained in terms of deficits to rule-based, language-specific structures (e.g., van der Lely, 2004). Versions of this theory include an impairment in specific structural relationships (agreement, specifier head-relations), absent linguistic features, fixation in a period of development where tense marking is ‘optional’, and problems in more general language functions (implicit rule learning, representing relationships between structures). Second, SLI has been explained in terms of a non-linguistic processing deficit that happens to particularly impact on language (e.g., Joanisse, in press). Proposals on the nature of this impairment include reduced processing rate, capacity limitations on cognitive processing, a deficit that particularly affects phonology, and a low-level perceptual or temporal processing deficit. Third, a recent neurobiological proposal by Ullman and Pierpont (2005) called the *Procedural-Declarative theory* argues that grammar acquisition is like skill learning, and therefore relies on procedural or implicit memory. By contrast, vocabulary acquisition concerns the learning of explicit knowledge and therefore relies on declarative memory. SLI corresponds to a developmental impairment of the procedural system. All of these

theories identify the deficits in SLI as involving disruptions to the language information stream in Figure 1.

Ullman and Pierpont's (2005) proposal is notable in that it identifies *compensation* as a key feature in producing the language profile of children with SLI. In the face of an impairment to the procedural learning system, Ullman and Pierpont argue that the declarative memory system attempts to compensate by acquiring certain aspects of language, such as frequently used phrases or inflected words. So, for example, where a typically developing child might inflect an English past tense such as 'talked' in terms of the regularities that operate in inflectional morphology (in English, to form the past tense, add -ed to the verb stem), the child with SLI might succeed in inflecting this high frequency verb by learning it as an unanalysed whole (note, however, that the performance of these children on inflection tasks is generally fairly poor). The evidence for this is that where normal children inflect regular verbs equally accurately irrespective of their frequency, children with SLI show frequency effects, inflecting high frequency regulars more accurately than low frequency (van der Lely & Ullman, 2001). Frequency effects are taken to be the hallmark of the operation of declarative memory.

What is important about Ullman and Pierpont's approach is that it emphasises the atypical learning process. Impaired behaviour is the outcome of development working under different constraints, rather than about focal damage to a component of a static system. Note that damage to a static system might be appropriate to explain a similar behavioural deficit when observed in a normal adult who has suffered brain damage. For example, individuals suffering Broca's aphasia after left anterior damage exhibit particular problems in processing grammar. However, focal damage in normal otherwise healthy children before the age of 5-7 does not produce SLI; it causes

language delay followed by recovery to within the normal range (see Bates & Roe, 2001, for a review). Interestingly, the effects of early damage are similar irrespective of side of damage while in adults, impairments in processing the structural aspects of language only occur after left-sided damage. In short, then, SLI must be viewed as an atypical developmental process.

What is problematic in Ullman and Pierpont's approach is that it highlights that we don't really know what the atypical developmental process looks like (Thomas, 2005b). How does compensation actually work? Why is it not fully successful, in which case the atypical process would not produce surface behavioural impairments? The implication is that compensatory processes are limited in some respect but unless these processes are specified in detail, such a hypothesis cannot be falsified, thereby making the theory untestable. Two recent methodologies have begun to make progress in specifying the nature of compensatory processes.

One of the methodologies is the use of computational models of development to provide formal, implemented simulations of the proposed atypical process (Thomas & Karmiloff-Smith, 2003). This approach begins by building a computational model of normal development for a particular aspect of language acquisition, such as learning to produce past tenses or to parse sentences. The normal developmental trajectory is the consequence both of the linguistic environment to which the system is exposed and its internal computational constraints, such as the nature of its representations and learning algorithm. Manipulations to the linguistic environment and internal computational constraints provide candidate hypotheses to explain atypical development, if those manipulations are able to deflect the normal trajectory so that it now characterises the pattern observed in a particular disorder.

In this way, Thomas (2005b) demonstrated how altering a computational property in a connectionist model of English past tense acquisition was sufficient to deflect development from the normal trajectory to the SLI profile. This property was the discriminability of the internal processing units (corresponding roughly to the signal-to-noise ratio of a neural processing system). This manipulation was notable for three reasons. First, the property was altered in a processing channel that was shared by both regular inflections (*talk-talked*) and irregular inflections (*drink-drank*), yet it affected regular inflections more seriously than irregulars. This is because good discriminability is necessary to learn the sharp category boundaries in internal representations that will depict rules or regularities. Changes to shared resources can therefore produce uneven deficits to the separate processes that use those resources. Second, changing the processing property at the start of development altered the way the system exploited the information available to it. In the normal system, phonological input was preferentially used to drive regular past tense formation while lexical-semantic (word-specific) information was preferentially utilised to drive irregular past tense formation. In the inefficient, slowly developing atypical system, there was a greater reliance on word-specific lexical-semantic information to drive *all* past tense formation. This led to the emergence of frequency effects in regular past tense formation observed empirically by van der Lely and Ullman (2001), and is in line with the proposal that all verbs are treated as exceptions in SLI. Third, the model captured SLI accuracy levels in children of around ten years of age. However, the atypical model was then run on to predict adult performance. The results suggested resolution of difficulties on highly practised items, but residual difficulties when the system came to extend its knowledge to novel cases (i.e., applying the rule). In other

words, externally, the system eventually seemed to compensate for highly practised items but internally it failed to normalise.

Using a similar approach, Thomas and Redington (2004) constructed a recurrent connectionist model of sentence processing to simulate the results of an experiment in which participants had to identify the agent and patient of a sentence (Dick et al., 2001). In this task, participants heard sentences that were either canonical (active: *The dog chases the cat*; subject cleft: *It is the dog that chases the cat*) or non-canonical (passive: *The cat is chased by the dog*; object cleft: *It is the cat that the dog chases*) and were required to make a binary choice as quickly as possible on which of two pictures (*dog, cat*) corresponded to the agent (*dog*). Dick et al. (2001) found that adults with acquired aphasia exhibited marked difficulties at identifying the agents of non-canonical sentences, that is, both passives and object clefts. When the trained ‘adult’ connectionist model was lesioned, it too exhibited this pattern of deficits. However, when the same model had its processing resources reduced prior to training to simulate a developmental disorder, it predicted that the deficits should be more marked for object cleft sentences than passives.

Let us consider why this should be the case. In the atypical model, the resource limitation reduced the ability of the connectionist network to learn information across sequences of words. Object cleft sentences are identified by a noun-noun sequence (*cat that the dog*) and so suffer from developmental limitations in sequence processing. However, passive sentences are also identified by lexical cues (past participle *chased* and preposition *by*); across learning, the network was able to learn to use these cues to identify this construction. Importantly, when Dick et al. (2004) extended their paradigm to typically developing children and children with SLI, the results supported the prediction of the model: performance on passives and

object clefts was closely related in adult aphasics, while in children with SLI, passive constructions were identified more accurately than object clefts.

These models demonstrate the benefit of implementation for making theories more explicit. Together, the models demonstrate: (1) how adaptive learning systems do the best they can with atypical properties they possess; (2) that compensated systems may use information sources in different ways; and (3) that atypical processing properties may allow compensation for some parts of language but not others.

A second methodology essential to uncover the nature of compensation in developmental disorders is that of functional brain imaging. The computational simulations suggest that, with age and practise, behavioural problems can resolve even though the underlying processes have not normalised. If so, behavioural measures, especially those with poor sensitivity such as standardised tests, may be insufficient to assess developmental outcome. By contrast, functional brain imaging offers a window on the way in which the brain has adapted to perform language tasks when its computational constraints are atypical.

Using this approach, we recently imaged the brain of a 42-year-old man called CK who was diagnosed with SLI aged 6 (Richardson et al., 2006). Donlan et al. (2006) compared the language profile of CK available from standardised tests and educational records when he joined a special school for children with language impairments in 1971, with his performance as an adult in order to explore the eventual outcome of language development. CK's school records indicated a verbal IQ of 69 at 6 years of age, and particular difficulties with auditory memory and morphological inflections. The records note that CK had reduced babbling as a baby, he used only 3 words used at two years of age (girl, pig, stop) and there was then no further

productive output until 5 years and 3 months (he started receiving speech and language therapy at 4 years and 11 months). CK's adult profile indicated that some aspects of his language were now within or above the normal range: receptive vocabulary was in the 99th percentile, auditory discrimination was at ceiling, picture comprehension was in the 63rd percentile, and naming showed a z-score of 0.16, i.e., slightly above average. However, CK revealed persisting deficits in tasks requiring phonological working memory: non-word repetition had a z-score of -1.94, well below the normal range, and recall of sentences as in the 1st percentile.

Functional imaging was used to explore brain activations in CK during passive listening to sentences, or reading of sentences presented one word at a time at the same rate, against a baseline of backwards speech or nonsense visual symbols. CK's performance was compared to a group of 14 adult controls. The results revealed that for CK, there was reduced activation in temporal regions normally associated with phonological processing, but *increased* activation in dorsal pre-motor and superior temporal regions, as well as in the caudate nucleus. The latter are all motor areas but note that the task CK was asked to perform included no motor component. One must interpret results of this form with care, since there are at least three ways one could explain the differences between CK and controls: (1) as adaptive compensation; (2) as a failure of the system to inhibit task-irrelevant circuits; (3) as a case of task-irrelevant activations causing interference (though those activations might be adaptive for some other task). Nevertheless, one possible interpretation of the findings is that CK was using additional sub-articulation during comprehension as a compensatory process to support semantic retrieval during language comprehension. Interestingly, Vargha-Khadem et al. (1998) also reported increased activation in the caudate nucleus in

language tasks in the affected members of the KE family. However, those individuals also showed increased activation in Broca's area, a pattern not observed in CK.

In sum, current research of developmental disorders of language is exploiting multiple, interdisciplinary methods, including genetic, computational, and brain imaging in an attempt to better characterise the nature of the atypical developmental process (see Mareschal et al., 2007, for a review of a similar multidisciplinary approach to developmental dyslexia).

5. Conclusion

Developmental disorders of language can exhibit contrasting profiles of strength and weakness. These can be traced to different information streams involved in the task of language learning. The relation of atypical language systems (such as those observed in Williams syndrome and Specific Language Impairment) to the normally developing system remains controversial, but perhaps the best approach is to view them as shedding light on the constraints that shape the learning process. However, the onus then moves onto specifying the detailed nature of this learning process, involving such ideas as redundancy (illustrated in the example of WS) and compensation (illustrated in the example of SLI). New methodologies such as computational modelling and functional brain imaging will be important complements to behavioural studies in this endeavour.

Acknowledgements

This work was supported by British Academy Grant SG – 40400 and UK Medical Research Council Grant G0300188 to Michael Thomas.

References

- Annaz, D., Johnson, M. H., & Thomas, M. S. C. (2006). Contrasting cognitive profiles of 5-12 year old children with Williams syndrome, Down syndrome, high-functioning autism, and low-functioning autism: A comparison of developmental trajectories. *Manuscript in preparation*.
- Bates, E., & Roe, K. (2001). Language development in children with unilateral brain injury. In C. A. Nelson & M. Luciana (Eds.), *Handbook of Developmental Cognitive Neuroscience* (p. 281-307). Cambridge, Mass: MIT Press.
- Bishop, D. V. M. (1997). Cognitive neuropsychology and developmental disorders: Uncomfortable bedfellows. *Quarterly Journal of Experimental Psychology*, 50A, 899-923.
- Bishop, D. V. M., & Norbury, C. F. (2002). Exploring the borderlands of autistic disorder and specific language impairment: A study using standardised diagnostic instruments. *Journal of Child Psychology and Psychiatry*, 43, 917-929.
- Brock, J. (in press). Language abilities in Williams syndrome: a critical review. *Development and Psychopathology*
- Chiat, S. (2001). Mapping theories of developmental language impairment: Premises, predictions and evidence. *Language and Cognitive Processes*, 16, 113-142.
- Clahsen, H., & Temple, C. (2003). Words and rules in Williams syndrome. In Y. Levy, Y. & J. Schaeffer (Eds.), *Towards a definition of Specific Language Impairment in children*. Erlbaum.
- Dick, F., Bates, E., Wulfeck, B., Aydelott, J., Dronkers, N., & Gernsbacher, M. (2001). Language deficits, localization, and grammar: Evidence for a distributive model of language breakdown in aphasic patients and neurologically intact individuals. *Psychological Review*, 108(3), 759-788.

- Dick, F., Wulfeck, B., Krupa-Kwiatkowski, & Bates (2004). The development of complex sentence interpretation in typically developing children compared with children with specific language impairments or early unilateral focal lesions. *Developmental Science*, 7(3), 360-377.
- Donlan, C., Aboagye, S., Clegg, J. & Stackhouse, J. (2006). Cognitive-developmental processes in individuals with Specific Language Impairments: Three cases observed in childhood and mid-life. *Manuscript in preparation*.
- Donnai, D., & Karmiloff-Smith, A. (2000) Williams Syndrome: from genotype through to the cognitive phenotype. *American Journal of Medical Genetics*, 97, 164-171.
- Dunn, Ll. M., Dunn, L.M., Whetton, C. & Burley, J. (1997). *British Picture Vocabulary Scale 2nd edition (BPVS-II)*. NFER-Nelson Publishing Company Limited: Windsor, Berks.
- Elliott, C. D., Smith, P. & McCulloch, K. (1996). *British Ability Scales Second Edition (BAS II)*. NFER-Nelson Publishing Company Limited: Windsor, Berks.
- Fowler, A. (1998). Language in mental retardation: Associations with and dissociations from general cognition. In J. A. Burack, R. M. Hodapp, & E. Zigler (Eds.), *Handbook of Mental Retardation and Development* (p. 290-333). Cambridge: Cambridge University Press.
- Goldin-Meadow, S. (2005). *The resilience of language*. Psychology Press: Hove, Sussex.
- Happé, F. (1994). *Autism*. UCL Press: London.
- Joanisse, M.F. (in press). Phonological deficits and developmental language impairments. In D. Mareschal, S. Sirois & G. Westermann (Eds.),

- Neuroconstructivism Volume II: Perspectives and Prospects*. Oxford University Press: Oxford.
- Jones, W., Bellugi, U., Lai, Z., Chiles, M., Reilly, J., Lincoln, A., & Adolphs, R. (2000). Hypersociability in Williams syndrome. *Journal of Cognitive Neuroscience, 12: Supplement*, 30-46.
- Karmiloff-Smith, A. (1998). Development itself is the key to understanding developmental disorders. *Trends in Cognitive Sciences, 2(10)*, 389-398.
- Laing, E., Butterworth, G., Ansari, D., Gsödl, M., Laing, E., Barnham, Z., Lakusta, L., Tyler, L.K., Grice, S., Paterson, S., & Karmiloff-Smith, A. (2002). Atypical linguistic and socio-communicative development in toddlers with Williams syndrome. *Developmental Science, 5 (2)*, 233-246.
- Marcus, G. F. & Fisher, S. E. (2003). FOXP2 in focus: what can genes tell us about speech and language? *Trends in Cognitive Sciences, 7(6)*, 257-262.
- Mareschal, D., Johnson, M., Sirios, S., Spratling, M., Thomas, M. S. C., & Westermann, G. (2007). *Neuroconstructivism Volume 1: How the brain constructs cognition*. Oxford: Oxford University Press.
- McDonald, J. L. (1997). Language acquisition: The acquisition of linguistic structure in normal and special populations. *Annual Review of Psychology, 48*, 215-241.
- Morton, J. (2004). *Developmental disorders: A causal modelling approach*. Blackwell Publishing: Oxford.
- Newbury, D. F., et al. (2002). FOXP2 is not a major susceptibility gene for autism or specific language impairment. *American Journal of Human Genetics, 70*, 1318-1327.
- Paterson, S. J., Brown, J. H., Gsödl, M. K., Johnson, M. H. & Karmiloff-Smith, A. (1999). Cognitive Modularity and Genetic Disorders. *Science, 286*, 2355-2358.

- Pinker, S. (1994). *The language instinct*. Penguin: London
- Pinker, S. (1999). *Words and rules*. Weidenfeld & Nicolson: London.
- Rice, M., Warren, S. F., & Betz, S. K. (2005). Language symptoms of developmental language disorders: An overview of autism, Down syndrome, fragile X, specific language impairment, and Williams syndrome. *Applied Psycholinguistics*, *26*, 7-27.
- Richardson, F., Thomas, M. S. C., Donlan, C., Crinion, J., & Price, C. (2006). Case study of a 42-year-old man who had SLI as a child: functional imaging of an atypical language system. *Manuscript in preparation*.
- Ring, M. & Clahsen, H. (2005). Distinct patterns of language impairment in Down's syndrome, Williams syndrome, and SLI: The case of syntactic chains. *Journal of Neurolinguistics*, *18*, 479-501.
- Tager-Flusberg, H. & Sullivan, K. (1998). Early language development in children with mental retardation. In E. J. Burack, R. Hodapp, & E. Zigler (Eds.), *Handbook of development and retardation* (pp. 208-239). New York: Cambridge University Press.
- Temple, C. (1997). *Developmental cognitive neuropsychology*. Psychology Press: Hove, Sussex.
- Thomas, M. S. C. (2005a). Constraints on language development: Insights from developmental disorders. In: P. Fletcher & J. Miller (Eds.), *Language disorders and developmental theory*, (p. 11-34). John Benjamins: Philadelphia.
- Thomas, M. S. C. (2005b). Characterising compensation. *Cortex*, *41*(3), 434-442.
- Thomas, M. S. C. (2006). Williams syndrome: Fractionations all the way down? *Cortex*.

- Thomas, M. S. C., Dockrell, J. E., Messer, D., Parmigiani, C., Ansari, D., & Karmiloff-Smith, A. (in press). Speeded naming, frequency and the development of the lexicon in Williams syndrome. *Language and Cognitive Processes*.
- Thomas, M. S. C. & Karmiloff-Smith, A. (2002). Are developmental disorders like cases of adult brain damage? Implications from connectionist modelling. *Behavioural and Brain Sciences*, 25(6), 727-780.
- Thomas, M. S. C. & Karmiloff-Smith, A. (2003). Modelling language acquisition in atypical phenotypes. *Psychological Review*, 110(4), 647-682.
- Thomas, M. S. C., & Karmiloff-Smith, A. (2005). Can developmental disorders reveal the component parts of the human language faculty? *Language Learning and Development*, 1(1), 65-92.
- Thomas, M. S. C. & Redington, M. (2004). Modelling atypical syntax processing. In W. Sakas (Ed.), *Proceedings of the First Workshop on Psycho-computational models of human language acquisition at the 20th International Conference on Computational Linguistics*. Pp. 85-92.
- Toga, A. W., Thompson, P. M., & Sowell, E. R. (2006). Mapping brain maturation. *Trends in Neurosciences*, 29(3), 148-159.
- Tomasello, M., & Farrar, M. J. (1986). Joint attention and early language. *Child Development*, 57, 1454-1463.
- Ullman, M. T. & Pierpont, E. I. (2005). Specific Language Impairment is not specific to language: The Procedural Deficit Hypothesis. *Cortex*, 41(3), 399-433.
- van der Lely, H. K. J. (2004). Evidence for and implications of a domain-specific grammatical deficit. In L. Jenkins (Ed.), *The genetics of language*, (p. 117-145). Elsevier, Oxford.

- van der Lely, H. K. J. & Ullman, M. T. (2001). Past tense morphology in specifically language impaired and normally developing children. *Language and Cognitive Processes*, 16, 177-217.
- Vargha-Khadem F., Watkins, K. E., Price C. J., Ashburner, J., Alcock, K., Connelly, A., Frackowiak, R. S. J., et al. (1998). Neural basis of an inherited speech and language disorder. *Proceedings of the National Academy of Sciences USA*, 95, 12695-12700
- Watkins, K. E., Dronkers, N. F., & Vargha-Khadem, F. (2002). Behavioural analysis of an inherited speech and language disorder: comparison with acquired aphasia. *Brain*, 125, 454-464.
- Watkins, K. E., Vargha-Khadem, F., Ashburner, J., Passingham, R. E., Friston, K. J., Connelly, A., Frackowiak, R. S. J., Mishkin, M. & Gadian, D.G. (2002). MRI analysis of an inherited speech and language disorder: structural brain abnormalities. *Brain*, 125, 465-478.

Figure Captions

Figure 1. Information streams combined in language acquisition, along with developmental disorders in which the primary deficits relate to one of the streams

Figure 2. Cross-sectional developmental trajectories for children with different developmental disorders on two standardised tests (Annaz, Johnson, & Thomas, 2006). Left panel: British Picture Vocabulary Scale (Dunn et al., 1997); right panel: Pattern Construction from the British Abilities Scales (Elliott et al., 1996). ASD = Autistic spectrum disorder, HF = high functioning, LF = low functioning, DS = Down syndrome, WS = Williams syndrome, TD = typically developing controls

Figure 3. Developmental fractionation of cognition in Williams syndrome: (a) early characterisation: genetic mutation produces simple fractionation between general cognition and language; (b) subsequent research indicates complex pattern of fractionation in both linguistic and non-linguistic domains (Thomas, 2006). Labelled boxes indicate dissociations reported by one or more studies in the literature. Triangles indicate domains in which there is a scale of difficulty and individuals with WS have been reported to show exaggerated deficits on harder parts of the domain.

Figure 1.




Figure 2




Figure 3a




Figure 3b

